

The Unique Derivative Expansion for Thermal Effective Actions*

T.S.Evans^{†‡}

Theoretical Physics, Blackett Laboratory, Imperial College, Prince Consort Road, London SW7 2BZ, U.K.
(21st August 1998, revised 16th September 1998.)

I show that there is a unique and well behaved derivative expansion of an effective action at finite temperature. The result is true for all formalisms including the popular Closed Time Path and Imaginary Time methods.

PACS: 11.10.Wx, 11.10.Lm. Preprint Imperial/TP/97-98/69

The effective action has been a powerful weapon in the arsenal of quantum field theory for many years [1–4]. In most cases, effective actions are still too difficult to work with. One way of extracting physically useful results is to try a further approximation, the derivative expansion. In particular, for an effective action of a scalar field, the lowest order term of such an expansion is the effective potential or free energy, a foremost tool in the study of the different phases of a theory.

However it has been known for some time [5] that at any non-zero temperature the derivative expansion is not well behaved in that there is no one, unique answer [5–17]. This then destroys the link between the Free energy and the potential that controls the dynamical evolution of a scalar field such as one has many pictures of phase transitions.

Here I will describe how the existing calculations incorrectly use equilibrium Green functions to calculate such effective actions. When considering the behaviour of slowly varying fields in a thermal bath, i.e. the system described by a derivative expansion of an effective action, then one finds that one must use different types of Green function. This is then seen to give a unique derivative expansion of the effective action.

The example I will consider has two fields, ϕ and η , whose dynamics is described by the Lagrangian

$$\mathcal{L}[\phi, \eta] = \frac{1}{2}\eta\Delta^{-1}\eta - \frac{1}{2}g\phi\eta^2 + \mathcal{L}[\phi] \quad (1)$$

One obtains an effective action for the ϕ field by integrating out the η field as follows

$$Z = \int D\phi D\eta \exp\{i \int d^4x \mathcal{L}[\phi, \eta]\} \quad (2)$$

$$= \int D\phi \exp\{iS_{\text{eff}}[\phi] + i \int d^4x \mathcal{L}[\phi]\} \\ S_{\text{eff}}[\phi] = \frac{i}{2}\text{Tr} \{\ln [1 - g\phi(x)\Delta(x, y)]\} \quad (3)$$

Here the integration is exact (a pure Gaussian integral in η) which is not always the case. Thus the effective action for ϕ is the pure ϕ classical part $\int d^4x \mathcal{L}[\phi]$ plus $S_{\text{eff}}[\phi]$. The $S_{\text{eff}}[\phi]$ contains *all* η fluctuations - both quantum and statistical.

Note that I have made no detailed specifications about the initial configuration or the subsequent evolution of the ϕ fields. All I have specified is that it interacts with an η field in a particular way, and I will be assuming that the η field is in equilibrium. The whole point of an effective action, in this case for ϕ , is that it describes the behaviour of just a *small* part of a system but it contains the effects of all fields. We are then free to manipulate the effective action in a manner appropriate to the physical problem. For instance I can treat $S_{\text{eff}}[\phi]$ as a description of a classical theory, or use it as the basis of a full, non-equilibrium quantum path integral.

In general the aim is that $S_{\text{eff}}[\phi]$ will produce a simpler description and allow for better (non-perturbative?) approximations. However the problem is that $S_{\text{eff}}[\phi]$ is still too complicated - it is non-local in $\phi(x)$ as I will note below. The solution is to use further approximations, namely first to expand the $\ln$ and then to use a derivative expansion. The latter will be the focus of this work.

The validity of such further approximations, and indeed the usefulness of integrating out one field first rather than another, must always be justified by the nature of the physical problem. For instance if the ϕ field has a mass much less than the mass of the η field then it makes sense to first integrate out the slow fluctuations of the η field. A standard example is the Euler-Heisenberg effective action for electromagnetic fields in a QED plasma where the ϕ field would become the photon field, and the η field the electron field [3,13]. Given that the recipe I use to obtain an effective action for the simple model (1) is often appropriate in genuine physical problems, I will use it on (1) to illustrate generic problems and my solution to them.

From (3), the next step is to expand the $\ln$ and I find

$$S_{\text{eff}}[\phi] = \frac{i}{2}\text{Tr} \{\ln [1 - g\phi(x)\Delta(x, y)]\} \quad (4)$$

$$= S_{\text{eff}}^{(1)} + S_{\text{eff}}^{(2)} + \dots \quad (5)$$

$$= -\frac{ig}{2}\text{Tr} \{\phi(x)\Delta(x, y)\} \quad (6)$$

*hep-ph/9808383. Based on a talk given at TFT98 - the 5th Workshop on Thermal Field Theory, Regensburg, Germany, 10-14 August, 1998. Original overheads available at <http://euclid.tp.ph.ic.ac.uk/~time/papers/reg/regtalk.ps>

[†]email: T.Evans@ic.ac.uk

[‡]WWW: <http://euclid.tp.ph.ic.ac.uk/~time>

$$-\frac{ig^2}{4}\text{Tr}\{\phi(x)\Delta(x,y)\phi(y)\Delta(y,x)\}+\dots \quad (7)$$

$$= \text{diagram 1} + \text{diagram 2} + \dots \quad (8)$$

Diagram 1: A dashed circle with a vertical line from the bottom to a point labeled $\phi(x)$. A label $\Delta(x,x)$ is next to the circle.

Diagram 2: A dashed circle with two vertical lines, one from the top to a point labeled $\phi(y)$ and one from the bottom to a point labeled $\phi(x)$. Labels $\Delta(x,y)$ are next to the top and bottom arcs.

where the dashed lines represent the η field propagator, Δ . I will study the second term in this series, $S_{\text{eff}}^{(2)}$,

$$S_{\text{eff}}^{(2)} = -\frac{ig^2}{4} \int d^4x \int d^4y \{\phi(x)\Delta(x,y)\phi(y)\Delta(y,x)\} \quad (9)$$

as this is the quadratic term in ϕ and so contains corrections to the mass and kinetic terms of ϕ .

$S_{\text{eff}}^{(2)}$ is still too complicated. As classical actions contain one space-time integral, I wish to keep one, say $\int d^4x$, in $S_{\text{eff}}^{(2)}$ of (9). We need therefore to do the second $\int d^4y$ of (9). However, we see that we need to know ϕ at both x (considered fixed) and y points and we are integrating over all y i.e. the effective action is non-local. This is too much information to expect. We are more likely to know a lot about the field at one point in time and/or space. It is therefore convenient in many problems to express the field at any point in time or space, y , in terms of the field and its derivatives at one given point x . Thus we use a Taylor expansion to expand $\phi(y)$ as an infinite series.

$$\phi(y) = \phi(x) + (y-x)^\mu \partial_\mu \phi(x) + \dots \quad (10)$$

$$= \exp\{i(y-x)^\mu p_\mu\} \phi(x) \quad (11)$$

where $p_\mu = -i\partial_\mu$. Then $S_{\text{eff}}^{(2)}$ of (9) is given in terms of an infinite number of x terms, $\phi(x)$, $\partial_\mu \phi(x)$ etc., the known η propagator $\Delta(x-y)$, and a series of simple $(y-x)_\mu$ polynomials. This is the derivative expansion. Taken as a whole nothing is gained, non-locality is replaced by the need to know an infinite number of derivatives. Progress is made by *truncating* this derivative expansion so that we need specify only a finite number of local terms in a further approximation to our full effective action.

Under the above approximations the result is valid only in certain physical situations. The truncation of the ln expansion in (7) means that we are restricted to weak coupling and/or weak fields. The truncation of derivative expansion means that we must consider fields varying *slowly* in time and space.

Putting this all together gives the expression

$$S_{\text{eff}}^{(2)} = -\frac{1}{2} \int d^4x \phi(x) B(-i\partial_\mu) \phi(x) \quad (12)$$

$$-iB(p) = \frac{(-ig)^2}{2} \int \frac{d^4k}{(2\pi)^4} i\Delta(k) i\Delta(k+p) \quad (13)$$

$$= \text{diagram 3} \quad (14)$$

Diagram 3: A dashed circle with two vertical lines, one from the top and one from the bottom, both ending in dots.

Comparing (11) with (12) shows that the derivative expansion is an expansion in powers of $p^\mu = (E, \vec{p})^\mu$. If, for exemplary purposes, I just focus on time/energy $E \equiv -i\partial/\partial\tau$ then

$$S_{\text{eff}}^{(2)} = \int d^4x \left[B(p=0) \phi(x)^2 + \left(\frac{\partial B(p)}{\partial E} \right)_{p=0} \phi(x) \frac{\partial}{\partial \tau} \phi(x) + \frac{1}{2} \left(\frac{\partial^2 B(p)}{\partial E^2} \right)_{p=0} \phi(x) \frac{\partial^2}{\partial \tau^2} \phi(x) + \dots \right] \quad (15)$$

i.e. the four-momentum expansion of B gives the derivative expansion of $S_{\text{eff}}^{(2)}$.

The analysis above is standard at zero temperature [1–3]. Formally there seems to be no problem at finite temperature, one can imagine that the same Feynman diagrams are required but one just uses finite temperature rules. If one takes the retarded bubble diagram, $B = B_{\text{ret}}$, then all thermal field formalisms give

$$B_{\text{ret}}(p_\mu = (E, \vec{p})) = -\frac{g^2}{2} \int \frac{d^3\vec{k}}{(2\pi)^3} \sum_{s_0, s_1 = \pm 1} \frac{s_0 s_1}{4\omega\Omega} \frac{1}{E + s_0\omega + s_1\Omega} \frac{(e^{\beta(s_0\omega + s_1\Omega)} - 1)}{(e^{\beta s_0\omega} - 1)(e^{\beta s_1\Omega} - 1)} \quad (16)$$

where

$$\omega = [\vec{k}^2 + m^2]^{1/2}, \quad \Omega = [(\vec{k} + \vec{p})^2 + m^2]^{1/2}. \quad (17)$$

However such equilibrium B 's do *not* have a unique momentum expansion at any non-zero temperature! This has been known for a long time, e.g. see Abrahams and Tsuneto [5]. I learned this from Fujimoto [7]. A careful calculation for the scalar retarded bubble gives [8,9,19]

$$\lim_{E, \vec{p} \rightarrow 0} B_{\text{ret}}(E, \vec{p}) = (\text{T}=0 \text{ part}) - \frac{g^2}{8\pi^2} \int_m^\infty d\omega N(\omega) \frac{k}{\omega^2 + v^2 \gamma^2 m^2} \quad (18)$$

$$\simeq -\frac{g^2}{16\pi} \frac{1}{1 + \gamma} \frac{T}{m} \quad (19)$$

where

$$N(E) = [\exp(\beta E) - 1]^{-1} \quad (20)$$

$$v = \lim_{E, \vec{p} \rightarrow 0} \frac{|\vec{p}|}{E}, \quad \gamma = (1 - v^2)^{-1/2} \quad (21)$$

Only in the case $v = \infty$ ($E \rightarrow 0$ before $\vec{p} \rightarrow 0$) is the usual contribution to the effective potential or Free energy recovered.

The solution comes when we realise that we may *not* use just the finite temperature Feynman rules in the zero temperature diagrams to obtain a finite temperature result in a general calculation including derivative

expansions of effective actions. Apart from anything else, a Feynman diagram does *not* represent a unique function at non-zero temperature there are many distinct types with significantly different values (retarded, time-ordered, thermal Wightman, ...) all of which all have their roles in different physical problems [20]. One must always consider what the physical problem is at finite temperature and let that tell you which mathematical Green function can be used to extract the relevant physics.

In the case of derivative expansions of effective actions, everyone uses the *equilibrium* Green functions where *all* fields are assumed to be in equilibrium and periodic. Usually retarded Green functions are used [11] though this is not always the case [15,17]. The question is *why are we using equilibrium Green functions?* A truncated derivative expansion of a periodic field is *not periodic*. Such approximations to field $\phi(y)$ are not equilibrium fields.

$$\begin{array}{ccccc} \phi(\tau) & = & \phi(0) & + & \tau \dot{\phi}(0) & + \dots \\ \text{periodic} & & \text{constant,} & & \text{linear} & \\ \text{in Im}(\tau) & & \text{periodic} & & \text{not periodic} & \end{array}$$

So you MUST NOT assume that external legs represent equilibrium fields !

The Green functions relevant to derivative expansions of effective actions are *not* generally equilibrium ones. Of course! We always wanted to look at how slowly varying fields (i.e. ones which are clearly not in equilibrium) behave in an equilibrium *background* - η is in equilibrium, ϕ is not.

So let us repeat calculation but

- We will work with time not energy in the Feynman integrals. This means that we can do *all* path-ordered thermal formalisms easily.
- The η field is in equilibrium and I use the following form of the propagator [18] valid for all path-ordered thermal field theory formalisms

$$\Delta(\tau, \tau'; \vec{k}) = \frac{-i}{2\pi} \int_{-\infty}^{\infty} dE e^{-iE(\tau-\tau')} [\theta_C(\tau, \tau') + N(E)] \rho(E, \omega) \quad (22)$$

where θ_C is a contour theta function, $N(E)$ is the Bose-Einstein distribution (20), and ρ is the spectral function. For the free scalar field η , ρ is of the form $(\pi/\omega) \cdot [\delta(E - \omega) - \delta(E + \omega)]$ with ω given in (17).

- The ϕ field is not assumed to be periodic or in equilibrium so $\exp\{\beta E\} \neq 1$ etc. when E is the energy or time derivative associated with a ϕ field. I only use (11) when doing the $\int dy$.

This gives

$$S_{\text{eff}}^{(2)} = -\frac{ig^2}{4} \text{Tr} \{ \phi(x) \Delta(x, y) \phi(y) \Delta(y, x) \} \quad (23)$$

$$= \Delta(x, y) \text{ (loop diagram with } \phi(x) \text{ and } \phi(y) \text{ external legs)} \quad (24)$$

$$= -\frac{1}{2} \int_{\tau_i}^{\tau_i - i\beta} d\tau \int \frac{d^3 \vec{p}}{(2\pi)^3} \phi(\tau, \vec{p}) \mathcal{B}(E, p; \tau - \tau_i) \phi(\tau, -\vec{p})$$

$$\begin{aligned} & \mathcal{B}(E, \vec{p}; \tau - \tau_i) \\ &= \frac{g^2}{2} \int \frac{d^3 \vec{k}}{(2\pi)^3} \sum_{s_0, s_1 = \pm 1} \frac{N(s_0 \omega) N(s_1 \Omega)}{4(s_0 \omega)(s_1 \Omega)} (F + G) \quad (25) \\ & \neq B_{\text{ret}} \end{aligned}$$

$$F = \frac{(e^{\beta A} - 1)}{A}, \quad (26)$$

$$G = \frac{(e^{-i(\tau - \tau_i)A} - 1)}{A} \cdot \frac{e^{\beta(s_0 \omega + s_1 \Omega)}}{N(E)} \quad (27)$$

$$A = E + s_0 \omega + s_1 \Omega \quad (28)$$

where $E = -i \frac{\partial}{\partial \tau}$ and acts only on the right hand side ϕ field. The τ_i is the start of the path in complex time used to define path-ordered thermal field formalisms [18]. It is the time at which the density matrix is given in closed-time path methods for instance. Since the derivative expansion requires an expansion of $\mathcal{B}$ about $E, \vec{p} \rightarrow 0$ then we are interested in the $\Omega \rightarrow \omega$ limit. When $s_0 = -s_1$ in the sum in (25) then $A \rightarrow 0$ and in this limit the denominators of F and G are zero. These $s_0 = -s_1$, $\omega - \Omega$ terms are produced by Landau damping processes.

This new result based on $\mathcal{B}$ has several consequences:-

1. $\mathcal{B}$ gives *unique* derivative series. When we take $E, p \rightarrow 0$ then the Landau damping terms have $A \rightarrow 0$. However both F and G are well behaved in this limit because of the form of the numerators.
2. In contrast, the equilibrium B_{ret} (16) is equivalent to setting $\exp\{\beta E\} = 1$ in $\mathcal{B}$ (25)¹, so that in B_{ret} $\exp\{A\} = \exp\{\pm \omega \pm \Omega\}$. Thus when we look at the Landau damping terms in B_{ret} where $A \rightarrow 0$ we find that $G \rightarrow 0$, but $F \sim A^{-1}$. This divergence in the integrand (there is still an integration over loop momenta to be done) leads to the poor behaviour of the four-momentum expansion of B_{ret} .
3. The only point where the correct $\mathcal{B}$ and old retarded equilibrium B_{ret} agree is in the *static* limit

¹One must also manipulate the pure ω dependent half of the integral in $\mathcal{B}$, switching variables to $\vec{k} + \vec{p}$ and integrating over this to get the form (16).

- $E = 0$. This is the only case where the derivative expansion of the field is in terms of a periodic field as a truly static i.e. time-independent, is trivially periodic. Another way of viewing this result is to say that this justifies the often ad-hoc rule used in some early effective potential/free energy calculations where the zero energy limit was taken first [11].

We can now write out the derivative expansion. Limiting myself to first order in derivatives then I find that for *any* path-ordered thermal field formalism the effective quadratic part of the effective Lagrangian $\mathcal{L}$ is

$$\begin{aligned}\mathcal{L}_{\text{eff}}[\phi] &= (M^2 + \delta M^2)\phi^2(\tau, \vec{x}) \\ &\quad - i\beta \left(\frac{1}{2}\delta M^2 + \mu(\tau - \tau_i) \right) \phi(\tau, \vec{x}) \frac{\partial}{\partial \tau} \phi(\tau, \vec{x}) \\ &\quad + \dots \\ \delta M^2 &= -\frac{g^2}{8\pi^2} \int_0^\infty dk \frac{N(\omega)}{\omega} \simeq -\frac{g^2}{16\pi} \frac{T}{M} \\ \mu(\tau) &= \frac{g^2}{32\pi^2} \int_0^\infty dk N^2(\omega) \frac{k^2}{\omega^3} \left[-2i\tau\omega e^{\beta\omega} \right. \\ &\quad \left. + \frac{1}{2}(e^{-2i\tau\omega} - 1)(1 + e^{2i(\tau-i\beta)\omega}) \right]\end{aligned}\quad (29)$$

I have assumed that the classical $\mathcal{L}[\phi]$ of (1) gives an $M^2\phi^2$ contribution to this order. The key point is that this is unique and independent of the order in which we take the zero energy, momentum/time, space derivative limit. It includes contributions from Landau Damping terms which cause problems in the zero momentum limits of equilibrium Green functions, yet there is no problem when using $\mathcal{B}$. In fact equilibrium retarded Green functions such as B_{ret} describe the response to a system to a delta function impulse, as this is what the usual linear response analysis tells us [18]. Thus such Green functions are relevant only for *sudden* changes. The derivative expansion of an effective action and $\mathcal{B}$ are appropriate for slowly varying fields, a very different type of physical problem.

This analysis of the derivative expansion of an effective action for scalar fields has used a general path ordered approach to thermal field theory using the two scalar field model of (1). The same model has also been considered in detail for the pure imaginary-time formalism [21] and in the closed-time path method [22,23]. The same result emerges in both cases as it must. However the result is viewed very differently in the two distinct approaches. Other more sophisticated models also have been considered - scalars or an abelian gauge field coupled to fermions in a heat bath, models with chemical potentials, and in effective actions for weakly coupled superconductors [24] - with similar success.

I would like to thank the organisers of TFT98 for creating such an exciting meeting where this talk was given. I would like to thank my collaborators [22–24] on various extensions of this work and Ray Rivers for continu-

ing discussions. Finally thanks to G.Aarts, I.Lawrie and C.van Weert for their perceptive comments on my talk at TFT98.

-
- [1] S.Coleman, *Aspects of Symmetry* (Cambridge University Press, Cambridge, 1985, ISBN 0 521 31827 0).
 - [2] C.M.Fraser, Z.Phys.C **28** 101 (1985).
 - [3] A.Dobado, A.Gómez-Nicola, A.L.Maroto, J.R.Peláez, *Effective Lagrangians for the Standard Model* (Springer-Verlag, Berlin, 1997, ISBN 3-540-62570-4).
 - [4] I.Moss, D.Toms and A.Wight, Phys.Rev.D **46** 1671 (1992).
 - [5] E.Abrahams and T.Tsuneto, Phys.Rev. **152** 416 (1966).
 - [6] Y.Fujimoto, *Puzzles in Finite Temperature Perturbation*, Kyoto Univ. preprint, RIFP-581, 1984, unpublished.
 - [7] Y.Fujimoto, in *Quantum Field Theory*, edited by F.Mancini (Elsevier Science Publishers, Amsterdam, 1986), p.435.
 - [8] T.S. Evans, Z.Phys.C **36** 153 (1987).
 - [9] T.S. Evans, Z.Phys.C **41** 333 (1988).
 - [10] Ch.G. van Weert, in *Proceedings of the 2nd Workshop on Thermal Field Theories and their Applications, Tsukuba, Japan, July 1990*, ed. H. Ezawa, T. Arimitsu, Y. Hashimoto, (Elsevier Sci. Pub., Amsterdam, 1991) p.17.
 - [11] T.S.Evans, Can.J.Phys. **71** 241 (1993).
 - [12] P.Elmfors, K.Enqvist, and I.Vilja, Phys.Rev.Lett. **71** 480 (1993), [hep-ph/9303269](#).
 - [13] P.S.Gribosky and B.R.Holstein, Z.Phys.C **47** 214 (1990).
 - [14] H.A.Weldon, Phys.Rev.D **47** (1993) 594.
 - [15] A.Das and M.Hott, Phys.Rev.D **50** 6655 (1994).
 - [16] I.J.R.Aitchison and D.J.Lee, Phys.Rev.B **56** 8303 (1997).
 - [17] A.Das, *Finite Temperature Field Theory* (World Scientific, Singapore, 1998).
 - [18] M.LeBellac, *Thermal Field Theory* (Cambridge University Press, Cambridge, 1996, ISBN 0 521 46040 9).
 - [19] T.S.Evans, *Thermal Bubble Diagrams Near Zero Energy*, Imperial/TP/92-93/45, [hep-ph/9307335](#), unpublished.
 - [20] T.S.Evans, Nucl.Phys. **B374** 340 (1992).
 - [21] T.S.Evans, *Derivative expansions of Euclidean thermal effective actions*, Imperial/TP/97-98/021, [hep-ph/9808382](#)
 - [22] M.Asprouli and V. Galan Gonzalez, *Derivative expansions of real-time thermal effective actions*, Imperial/TP/97-98/026, [hep-ph/9802319](#).
 - [23] M.Asprouli, *A way to get a well-defined derivative expansion of real-time thermal effective actions*, — Talk given at TFT98 - the 5th Workshop on Thermal Field Theories, Regensburg, Germany, August 10-14th August 1998, Imperial/TP/97-98/078, [hep-th/9809099](#).
 - [24] T.S.Evans and D.A.Steer, *Time-dependent effective theory of a weakly coupled superconductor at finite temperature*, in preparation.